



The Ragdoll is a semi-longhaired cat breed invented in the US during the 1960s and introduced to the UK in the 1980s. The Ragdoll name originates from their typically placid, friendly nature and tendency to go limp 'like a ragdoll' when picked up. However, although immensely popular worldwide, there has been limited published evidence on the overall health of the breed until now.

This study aimed to report on the demographics, common disorders, and mortality of the general population of Ragdoll cats under primary veterinary care in the UK in 2019.

Demography

included

1,254,484 cats
under UK vet
care during
2019.

The study


Ragdolls
comprised
21,358 (1.70%)
of the overall cat
population.


Average adult
Ragdoll bodyweight
was 4.46kg.
Males (4.97kg) were
heavier than
females (3.83kg)

during 2005-2019, from 0.70% of all cats born in 2005 to 3.72% in 2019.

Annual proportional birth rates showed an increasing Ragdoll popularity

Most commonly diagnosed disorders in Ragdolls





Most common causes of death in Ragdolls


complications and ear discharge.

Males had higher probability of obesity.

The **median age at death** in Ragdolls overall was **12.85 years**. Life expectancy did not differ statistically between females and males.

Kidney disorder 21.18%

Of the deaths in the study, 91.59% involved euthanasia and 8.41% died



Given its high frequency and impact on welfare, periodontal

(dental) disease should be considered a priority welfare issue for Ragdoll cats. Owners should seek veterinary advice to ensure good dental health in their Ragdoll cats.



Vet Compass



Veterinary College The current study reports the Ragdoll as a relatively common purebred cat breed in the UK and with steeply rising levels of ownership. Overall, the Ragdoll cat breed appears to be largely typical in terms of disorders and mortality for purebred cats overall in the UK.

CLICK HERE TO READ THE FULL STUDY